

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):


- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



PCT

WORLD INTELLECTUAL PRO  
International

INTERNATIONAL APPLICATION PUBLISHED UNDER

WO 9605823A1

(51) International Patent Classification 6 :

A61K 31/38

A1

(11) International Publication Number:

WO 96/05823

(43) International Publication Date: 29 February 1996 (29.02.96)

(21) International Application Number: PCT/US95/10581

(22) International Filing Date: 21 August 1995 (21.08.95)

(30) Priority Data: 08/294,238 22 August 1994 (22.08.94) US

(71) Applicant: ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).

(72) Inventor: GLASEBROOK, Andrew, Lawrence; 11410 Valley Meadow Drive, Zionsville, IN 46077 (US).

(74) Agent: LAMMERT, Steven, R.; Barnes &amp; Thornburg, 1313 Merchants Bank Building, Indianapolis, IN 46204 (US).

(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).



Published

With international search report.

(54) Title: METHODS OF INHIBITING DEMYELINATING AND DYSMYELINATING DISEASES

## (57) Abstract

A method of inhibiting demyelinating or dysmyelinating diseases or their symptoms comprising administering to a human in need thereof an effective amount of a compound having formula (I) wherein  $R^1$  and  $R^3$  are independently hydrogen,  $-CH_3$ , (a) or (b), wherein Ar is optionally substituted phenyl;  $R^2$  is selected from the group consisting of pyrrolidine, hexamethyleneimino, and piperidino; or a pharmaceutically acceptable salt of solvate thereof.



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

-1-

## METHODS OF INHIBITING DEMYELINATING AND DESMYELINATING DISEASES

Diseases of the myelin sheath are divided into two categories. The first category is the demyelinating or myelinoclastic diseases and the second is the dysmyelinating diseases. The myelinoclastic (demyelinating) diseases contain multiple sclerosis, myelinoclastic diffuse sclerosis, post-infectious and postvaccinal encephalomyelitis (disseminated vasculomyelinopathy), transverse myelitis, central pontine myelinolysis, and marchiafava-bignami disease.

The dysmyelinating diseases (leukodystrophies) contain metachromatic leukodystrophy, globoid cell leukodystrophy, adrenoleukodystrophy, and spongy sclerosis.

The first group (the demyelinating diseases) includes conditions in which the myelin sheath has developed normally and has a completely normal metabolic maintenance system, but in which the sheath appears to be the primary target of various conditions. In this group, two subgroups are recognized. The first one is multiple sclerosis, and some of its variants. The second group consists of the complications of various infections, principally viral, and vaccinations, which result from a misdirection of the immune response that has been activated by the infection or vaccination. Therefore, both humoral and cellular (delayed) immune factors cause the myelin sheath of either the central or the peripheral nervous systems, or of both, to become inflamed, edematous, or destroyed.

In the second category, the dysmyelinating diseases, an inborn error of metabolism causes a disturbance of myelinogenesis. The dysmyelination may result from a metabolic failure of the myelin maintenance system after normally formed myelin has been laid down.

The dysmyelinating diseases are of many types and include such disparate conditions as the leukodystrophies the gangliosidoses, such as Tay-Sachs disease; several amino-acidopathies, such as phenylketonuria and maple syrup urine disease; and probably other metabolic disturbances which in one way or another interfere with the normal development of myelin.


It should be pointed out that destruction of myelin is a nonspecific result of almost any injury to the nervous system and can be seen following physical trauma or vascular insults, with infections and neoplasms, or as a result of various intoxications. The conditions considered here are those in which such causes and factors are not recognized. Two conditions, central pontine myelinolysis and Marchiafava-Bignami disease, have been included because they fulfill the criteria of being primary diseases of myelin, although they are not easily classifiable as either variants of multiple sclerosis or instances of postinfectious or postvaccinal reactions of the nervous system.

The conditions to be described here affect primarily, but not exclusively, the central nervous system. While multiple sclerosis is a disease restricted to the central nervous system, the leukodystrophies normally also involve the peripheral nerves; the postinfectious or postvaccinal reactions may be restricted to the central nervous system (encephalomyelitis), or peripheral nervous system (Guillain-Barré syndrome) or may involve both to varying degrees.

In the experimental allergic encephalomyelitis (EAE) animal model, administration of myelin basic protein (MBP) induces EAE (Higgins et al., *J. Immunol.*, 140(2), 440-445, January 15, 1988; Bitar et al., *Cell. Immun.*, 112, 364-370, (1988), and is characterized by increased levels of TGF $\beta$  and IL-5 in the brain (Khoury et al., *J. Exp. Med.*, 176, 1355-1364, Nov. 1992). Agents which induce TGF $\beta$  and

other antiinflammatory cytokine(s) may be useful for the described diseases and their symptoms.

5 This invention provides methods of inhibiting demyelinating or dysmyelinating diseases or their symptoms comprising administering to a human in need thereof an effective amount of a compound of formula I



wherein R¹ and R³ are independently hydrogen, -CH₃,  $\text{-C(=O)-(C}_1\text{-C}_6\text{ alkyl)}$ , or  $\text{-C(=O)-Ar}$ , wherein Ar is optionally substituted phenyl;

15 R² is selected from the group consisting of pyrrolidino, hexamethyleneimino, and piperidino; and pharmaceutically acceptable salts and solvates thereof.

20 The current invention concerns the discovery that a select group of 2-phenyl-3-arylbenzothiophenes (benzothiophenes), those of formula I, are useful for inhibiting demyelinating or dysmyelinating diseases or their symptoms. The compounds of formula I induce the expression of TGFβ, and more specifically TGFβ-3, and this

25 indicates usefulness for inhibition of the diseases or their symptoms.

5 The therapeutic and prophylactic treatments provided by this invention are practiced by administering to a human in need thereof a dose of a compound of formula I or a pharmaceutically acceptable salt or solvate thereof, that is effective to inhibit demyelinating or dysmyelinating diseases or their symptoms or its symptoms.

10 The term "inhibit" includes its generally accepted meaning which includes prohibiting, preventing, restraining, and slowing, stopping or reversing progression, severity or a resultant symptom. As such, the present method includes both medical therapeutic and/or prophylactic administration, as appropriate.

15 Raloxifene is a preferred compound of this invention and it is the hydrochloride salt of a compound of formula 1 wherein  $R^1$  and  $R^3$  are hydrogen and  $R^2$  is 1-piperidinyl.

20 Generally, at least one compound of formula I is formulated with common excipients, diluents or carriers, and compressed into tablets, or formulated as elixirs or solutions for convenient oral administration, or administered by the intramuscular or intravenous routes. The compounds can be administered transdermally, and may be formulated as sustained release dosage forms and the like.

25 The compounds used in the methods of the current invention can be made according to established procedures, such as those detailed in U.S. Patent Nos. 4,133,814, 4,418,068, and 4,380,635 all of which are incorporated by reference herein. In general, the process starts with a benzo[b]thiophene having a 6-hydroxyl group and a 2-(4-hydroxyphenyl) group. The starting compound is protected, acylated, and deprotected to form the formula I compounds. Examples of the preparation of such compounds are provided in the U.S. patents discussed above. The term "optionally substituted phenyl" includes phenyl and phenyl substituted  
30 once or twice with  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_4$  alkoxy, hydroxy, nitro, chloro, fluoro, or tri(chloro or fluoro)methyl.  
35



The compounds used in the methods of this invention form pharmaceutically acceptable acid and base addition salts with a wide variety of organic and inorganic acids and bases and include the physiologically acceptable salts which are often used in pharmaceutical chemistry. Such salts are also part of this invention. Typical inorganic acids used to form such salts include hydrochloric, hydrobromic, hydroiodic, nitric, sulfuric, phosphoric, hypophosphoric and the like. Salts derived from organic acids, such as aliphatic mono and dicarboxylic acids, phenyl substituted alkanolic acids, hydroxyalkanoic and hydroxyalkandioic acids, aromatic acids, aliphatic and aromatic sulfonic acids, may also be used. Such pharmaceutically acceptable salts thus include acetate, phenylacetate, trifluoroacetate, acrylate, ascorbate, benzoate, chlorobenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, methylbenzoate, o-acetoxybenzoate, naphthalene-2-benzoate, bromide, isobutyrate, phenylbutyrate,  $\beta$ -hydroxybutyrate, butyne-1,4-dioate, hexyne-1,4-dioate, caprate, caprylate, chloride, cinnamate, citrate, formate, fumarate, glycollate, heptanoate, hippurate, lactate, malate, maleate, hydroxymaleate, malonate, mandelate, mesylate, nicotinate, isonicotinate, nitrate, oxalate, phthalate, teraphthalate, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, propiolate, propionate, phenylpropionate, salicylate, sebacate, succinate, suberate, sulfate, bisulfate, pyrosulfate, sulfite, bisulfite, sulfonate, benzene-sulfonate, p-bromophenylsulfonate, chlorobenzenesulfonate, ethanesulfonate, 2-hydroxyethanesulfonate, methanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, p-toluenesulfonate, xylenesulfonate, tartarate, and the like. A preferred salt is the hydrochloride salt.

The pharmaceutically acceptable acid addition salts are typically formed by reacting a compound of

formula I with an equimolar or excess amount of acid. The reactants are generally combined in a mutual solvent such as diethyl ether or benzene. The salt normally precipitates out of solution within about one hour to 10 days and can be isolated by filtration or the solvent can be stripped off by conventional means.

Bases commonly used for formation of salts include ammonium hydroxide and alkali and alkaline earth metal hydroxides, carbonates, as well as aliphatic and primary, secondary and tertiary amines, aliphatic diamines. Bases especially useful in the preparation of addition salts include ammonium hydroxide, potassium carbonate, methylamine, diethylamine, ethylene diamine and cyclohexylamine.

The pharmaceutically acceptable salts generally have enhanced solubility characteristics compared to the compound from which they are derived, and thus are often more amenable to formulation as liquids or emulsions.

Pharmaceutical formulations can be prepared by procedures known in the art. For example, the compounds can be formulated with common excipients, diluents, or carriers, and formed into tablets, capsules, suspensions, powders, and the like. Examples of excipients, diluents, and carriers that are suitable for such formulations include the following: fillers and extenders such as starch, sugars, mannitol, and silicic derivatives; binding agents such as carboxymethyl cellulose and other cellulose derivatives, alginates, gelatin, and polyvinyl pyrrolidone; moisturizing agents such as glycerol; disintegrating agents such as calcium carbonate and sodium bicarbonate; agents for retarding dissolution such as paraffin; resorption accelerators such as quaternary ammonium compounds; surface active agents such as cetyl alcohol, glycerol monostearate; adsorptive carriers such as kaolin and bentonite; and lubricants such as talc, calcium and magnesium stearate, and solid polyethyl glycols.

-7-

The compounds can also be formulated as elixirs or solutions for convenient oral administration or as solutions appropriate for parenteral administration, for instance by intramuscular, subcutaneous or intravenous routes. Additionally, the compounds are well suited to formulation as sustained release dosage forms and the like. The formulations can be so constituted that they release the active ingredient only or preferably in a particular part of the intestinal tract, possibly over a period of time. The coatings, envelopes, and protective matrices may be made, for example, from polymeric substances or waxes.

The particular dosage of a compound of formula I required to inhibit demyelinating or dysmyelinating diseases or their symptoms according to this invention, will depend upon the severity of the condition, the route of administration, and related factors that will be decided by the attending physician. Generally, accepted and effective daily doses will be from about 0.1 to about 1000 mg/day, and more typically from about 50 to about 200 mg/day. Such dosages will be administered to a subject in need thereof from once to about three times each day, or more often as needed, and for a duration to effectively inhibit the disease(s) or symptom(s).

It is usually preferred to administer a compound of formula I in the form of an acid addition salt, as is customary in the administration of pharmaceuticals bearing a basic group, such as the piperidino ring. For such purposes the following oral dosage forms are available.

### Formulations

In the formulations which follow, "Active ingredient" means a compound of formula I.

Formulation 1: Gelatin Capsules

Hard gelatin capsules are prepared using the following:

Ingredient	Quantity (mg/capsule)
Active ingredient	0.1 - 1000
Starch, NF	0 - 650
Starch flowable powder	0 - 650
Silicone fluid 350 centistokes	0 - 15

The ingredients are blended, passed through a No. 45 mesh U.S. sieve, and filled into hard gelatin capsules.

Examples of specific capsule formulations of raloxifene that have been made include those shown below:

Formulation 2: Raloxifene capsule

Ingredient	Quantity (mg/capsule)
Raloxifene	1
Starch, NF	112
Starch flowable powder	225.3
Silicone fluid 350 centistokes	1.7

Formulation 3: Raloxifene capsule

Ingredient	Quantity (mg/capsule)
Raloxifene	5
Starch, NF	108
Starch flowable powder	225.3
Silicone fluid 350 centistokes	1.7

-9-

Formulation 4: Raloxifene capsule

<u>Ingredient</u>	<u>Quantity (mg/capsule)</u>
Raloxifene	10
Starch, NF	103
Starch flowable powder	225.3
Silicone fluid 350 centistokes	1.7

Formulation 5: Raloxifene capsule

5

<u>Ingredient</u>	<u>Quantity (mg/capsule)</u>
Raloxifene	50
Starch, NF	150
Starch flowable powder	397
Silicone fluid 350 centistokes	3.0

The specific formulations above may be changed in compliance with the reasonable variations provided.

10 A tablet formulation is prepared using the ingredients below:

Formulation 6: Tablets

<u>Ingredient</u>	<u>Quantity (mg/tablet)</u>
Active ingredient	0.1 - 1000
Cellulose, microcrystalline	0 - 650
Silicon dioxide, fumed	0 - 650
Stearate acid	0 - 15

15 The components are blended and compressed to form tablets.

Alternatively, tablets each containing 0.1 - 1000 mg of Active ingredient are made up as follows:

-10-

Formulation 7: Tablets

Ingredient	Quantity (mg/tablet)
Active ingredient	0.1 - 1000
Starch	45
Cellulose, microcrystalline	35
Polyvinylpyrrolidone (as 10% solution in water)	4
Sodium carboxymethyl cellulose	4.5
Magnesium stearate	0.5
Talc	1

5 The Active ingredient, starch, and cellulose are  
passed through a No. 45 mesh U.S. sieve and mixed  
thoroughly. The solution of polyvinylpyrrolidone is mixed  
with the resultant powders which are then passed through a  
No. 14 mesh U.S. sieve. The granules so produced are dried  
at 50°-60° C and passed through a No. 18 mesh U.S. sieve.  
10 The sodium carboxymethyl starch, magnesium stearate, and  
talc, previously passed through a No. 60 U.S. sieve, are  
then added to the granules which, after mixing, are  
compressed on a tablet machine to yield tablets.

15 Suspensions each containing 0.1 - 1000 mg of  
Active ingredient per 5 mL dose are made as follows:

Formulation 8: Suspensions

Ingredient	Quantity (mg/5-ml)
Active ingredient	0.1 - 1000 mg
Sodium carboxymethyl cellulose	50 mg
Syrup	1.25 mg
Benzoic acid solution	0.10 mL
Flavor	q.v.
Color	q.v.
Purified water to	5 mL

-11-

The Active ingredient is passed through a No. 45 mesh U.S. sieve and mixed with the sodium carboxymethyl cellulose and syrup to form a smooth paste. The benzoic acid solution, flavor, and color are diluted with some of the water and added, with stirring. Sufficient water is then added to produce the required volume.

### ASSAYS

#### Assay 1

Experimental allergic encephalomyelitis (EAE) is an acute inflammatory autoimmune disease of the central nervous system that can be elicited in rodents and is the major animal model for the study of multiple sclerosis (MS). EAE is a systemically initiated autoimmune disease in which spinal cord homogenate or myelin basic protein (MBP) prepared in suitable adjuvants, such as complete Freund's adjuvant (CFA), is injected to activate the peripheral immune system. Rapid migration of activated T cells from peripheral blood to the CNS then occurs where they initiate a localized inflammation and subsequent demyelination. A monophasic, acute or spontaneous relapsing-remitting chronic form of the disease may develop according to the mode of sensitization, genetic background and age of the animal. Alternatively, the chronic relapsing form of EAE can also be induced by injection of MBP-specific T cell lines or clones of the helper/inducer phenotype (CD4+). The resulting demyelination closely resembles that obtained by injection of MBP.

Typically, spinal cord homogenate or MBP is mixed with CFA at a 1:1 ratio. Rodents are injected s.c. in the posterior flank at multiple sites with not less than 1 mg sensitizing protein in a final volume of approximately 100 ul. Subsequent inoculations, similar to the primary, can be administered at 7 day intervals. Rodents usually demonstrate severe neurological symptoms as early as 13 days after primary inoculation with peak incidence levels

-12-

of EAE being reached by 21 days. Subcutaneous injection of as few as  $10^5$  MBP-specific T cell line or clone cells of the CD4+ helper/inducer phenotype is also effective at inducing EAE. Compound is administered orally or s.c., before or after primary antigen inoculation, and the beneficial effects determined by evaluation of neurological symptoms associated with EAE disease progression. One can also examine cerebrospinal fluid or brain for the presence of inflammatory cells, oligoclonal IgG or increased class II expression.

#### ASSAY 2

Five to fifty patients are selected for the clinical study. The patients suffer from a demyelinating or desmyelinating disease which exhibits symptoms, but otherwise are in good general health. Because of the idiosyncratic and subjective nature of these disorders, the study has a placebo control group, i.e., the patients are divided into two groups, one of which receives a compound of formula 1 as the active agent and the other receives a placebo. Patients in the test group receive between 50-200 mg of the drug per day by the oral route. They continue this therapy for 3-12 months. Accurate records are kept as to the number and severity of the symptoms in both groups and at the end of the study these results are compared. The results are compared both between members of each group and also the results for each patient are compared to the symptoms reported by each patient before the study began.


Utility of the compounds of formula I is illustrated by the positive impact they have in at least one of the assays described above.



-13-

I claim:

1. A method of inhibiting demyelinating or  
dysmyelinating diseases or their symptoms comprising  
5 administering to a human in need thereof an effective amount  
of a compound having the formula



(I)

wherein  $R^1$  and  $R^3$  are independently hydrogen,  
-CH<sub>3</sub>,  $\text{-}\overset{\text{O}}{\parallel}\text{C}\text{-(C}_1\text{-C}_6\text{ alkyl)}$ , or  $\text{-}\overset{\text{O}}{\parallel}\text{C-Ar}$ , wherein Ar is optionally  
substituted phenyl;


15  $R^2$  is selected from the group consisting of  
pyrrolidine, hexamethyleneimino, and piperidino; or a  
pharmaceutically acceptable salt of solvate thereof.

2. The method of Claim 1 wherein said compound is  
20 the hydrochloride salt thereof.

3. The method of Claim 1 wherein said  
administration is prophylactic.

-14-

4. The method of Claim 1 wherein said compound is



or its hydrochloride salt.

5. The method of Claim 1 wherein said disease is multiple sclerosis

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US95/10581

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 31/38

US CL : 514/443

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/443

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS

SEARCH TERMS: MULTIPLE SCLEROSIS, ENCEPHALOMYELITIS AND AROYLBENZOTHIOPHENE?

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US, A, 4,133,814 (JONES ET AL) 09 January 1979, Column 1, Line 10 to Column 2, Line 3.	1-5
A	US, A, 4,418,068 (JONES) 29 November 1983, Column 1, Line 10 to Column 2, Line 47.	1-5

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	T	later document published after the international filing date or priority date and not in conflict with the application but cited to understate the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, each such combination being obvious to a person skilled in the art
*L* document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	Z*	document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means		
*P* document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

26 SEPTEMBER 1995

Date of mailing of the international search report

19 OCT 1995

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer  
THEODORE J. CRIARES

Telephone No. (703) 308-1235

Form PCT/ISA/210 (second sheet)(July 1992)\*

THIS PAGE BLANK (USPTO)